



**Bayer CropScience**

Bayer CropScience LP  
 P.O. Box 12014, 2 T.W. Alexander Drive  
 Research Triangle Park, North Carolina 27709  
 1-866-99BAYER (1-866-992-2937)  
<http://www.bayercropscienceus.com>

**STRATEGO® Fungicide**

**For Use on Soybean to control Rust  
 (*Phakopsora pachyrhizi*)**

**SECTION 18 QUARANTINE EXEMPTION  
 File Symbol: 04-MN-05**

**FOR DISTRIBUTION AND USE ONLY IN THE STATE OF MINNESOTA  
 UNDER SECTION 18 QUARANTINE EXEMPTION**

This labeling must be in possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire Directions for Use and Disclaimer of Warranties before using this product. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for STRATEGO® Fungicide (EPA Registration Number 264-779).

Any adverse effects resulting from the use of STRATEGO® Fungicide under this quarantine exemption must be immediately reported to the State of Minnesota Department of Agriculture.

**EFFECTIVE PERIOD:**

This labeling becomes effective 12/13/04 and expires on 12/01/07.

A maximum of 2 total applications using approved Section 18 products collectively will be made under this soybean rust Section 18.

**DIRECTIONS FOR USE**

A maximum of 3.6 million acres of soybeans may be treated under this exemption.

**RECOMMENDED APPLICATIONS**

CROP	DISEASES	RATE OF STRATEGO® FUNGICIDE
Soybean	Rust ( <i>Phakopsora pachyrhizi</i> )	5.5 to 10 fl oz per acre

**Use Directions:** Begin broadcast, foliar applications of STRATEGO® preventatively when conditions are favorable for disease development. Alternate every application of STRATEGO® with at least one application of another effective (non-strobilurin) mode of action fungicide. Continue application programs as needed for disease control using a 10 to 14-day spray interval. Applications of STRATEGO® may be made up to the R6 (full seed) soybean growth stage. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. Do not apply STRATEGO® in combination with organosilicate surfactants or crop injury may occur. STRATEGO® should be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment. Chemical-resistant gloves must be worn for mixers/loaders. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

**Restrictions:** Applications may not be made within 21 days of harvest. Do not apply more than 2 applications per season. Do not apply more than 20 fl. oz/A per use season. Do not enter treated fields for 24 hours after application. Do not feed soybean forage or hay to livestock.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

In order to protect federally listed endangered and threatened species, the Minnesota Department of Agriculture has a program to pre-notify landowners where pesticide applications may affect federally listed endangered and threatened species.

However, no federally listed threatened or endangered species in Minnesota are likely to be adversely affected by these uses of STRATEGO® Fungicide.

**For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577**

**For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)**

or

**Visit our worldwide web site at [www.bayercropscienceus.com](http://www.bayercropscienceus.com)**

*As with any crop-protection product, always read and follow label directions.  
 For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*